

PATENT
3523/2/US

UNITED STATES UTILITY PATENT APPLICATION

for

**SOLUTION COMPOSITION OF AN OXAZOLIDINONE ANTIBIOTIC
DRUG HAVING ENHANCED DRUG LOADING**

Inventor:

Sandra M. Sims

EXPRESS MAIL MAILING LABEL

NUMBER EL 87720668845

DATE OF DEPOSIT August 20, 2001

I hereby certify that this Patent Application is being deposited with the United States Postal Service "EXPRESS MAIL POST OFFICE TO ADDRESSEE" service under 37 C.F.R. §1.10 on the date indicated above and is addressed to:

Assistant Commissioner for Patents, Box Patent Application, Washington, DC 20231.

Signature

Bernadette Y. Harper

SOLUTION COMPOSITION OF AN OXAZOLIDINONE ANTIBIOTIC DRUG
HAVING ENHANCED DRUG LOADING

This application claims the benefit of U.S. Provisional Application No.
5 60/285,347, filed April 10, 2001 and of U.S. Provisional Application No. 60/226,846,
filed 22 August 2000.

FIELD OF THE INVENTION

The present invention relates to a pharmaceutical composition in aqueous
solution form useful for oral, parenteral or topical application to a subject for
10 treatment or prevention of infective disease. In particular, the present invention
relates to such a composition having as an active agent an oxazolidinone antibiotic
drug. The field of the present invention also includes therapeutic or prophylactic use
of such a composition, and use of such a composition in preparation of a medicament.

BACKGROUND OF THE INVENTION

15 Numerous oxazolidinone compounds have been reported having
therapeutically and/or prophylactically useful antibiotic or antimicrobial, in particular
antibacterial, effect. Among such compounds are those illustratively disclosed in the
following patents, each of which is individually incorporated herein by reference.

U.S. Patent No. 5,164,510 to Brickner.

20 U.S. Patent No. 5,231,188 to Brickner.

U.S. Patent No. 5,565,571 to Barbachyn & Brickner.

U.S. Patent No. 5,627,181 to Riedl *et al.*


U.S. Patent No. 5,652,238 to Barbachyn *et al.*

U.S. Patent No. 5,688,792 to Barbachyn *et al.*

25 U.S. Patent No. 5,698,574 to Riedl *et al.*

U.S. Patent No. 6,069,145 to Betts.

Compounds disclosed in above-cited U.S. Patent No. 5,688,792 include for
example the compound (S)-N-[[3-[3-fluoro-4-(4-morpholinyl)phenyl]-2-oxo-5-
oxazolidinyl]methyl]acetamide, also referred to herein as linezolid. Linezolid has the
30 structure shown in formula (I):



and is in commercial use as a medicament under the trademark Zyvox® of Pharmacia Corporation. Linezolid exhibits strong antibacterial activity against gram-positive organisms including those of the following genera: *Staphylococcus* (e.g.,

- 5 *Staphylococcus aureus*, *Staphylococcus epidermidis*), *Streptococcus* (e.g., *Streptococcus viridans*, *Streptococcus pneumoniae*), *Enterococcus* (e.g., *Enterococcus faecalis*, *Enterococcus faecium*), *Bacillus*, *Corynebacterium*, *Chlamydia* and *Neisseria*. Many such gram-positive organisms have developed significant levels of resistance to other antibiotics. Oxazolidinone antibiotics are also generally effective against
- 10 anaerobic organisms such as those of the genera *Bacteroides* and *Clostridia*, and against acid-fast organisms such as those of the genus *Mycobacterium*.

Above-cited U.S. Patent No. 5,688,792 discloses that the subject antibiotic oxazolidinone compounds, including linezolid, can be formulated as liquid form compositions including solutions. For example, it is disclosed therein that a solution

15 can be provided of a subject oxazolidinone compound in water or in a water-propylene glycol or water-polyethylene glycol system, optionally containing conventional coloring agents, flavoring agents, stabilizers and thickening agents.

It is further disclosed in above-cited U.S. Patent No. 5,688,792 that the subject oxazolidinone compounds can be administered orally, parenterally and/or topically,

20 and that parenteral administration can be by intravenous injection or other parenteral route. For parenteral administration, it is disclosed therein that a suitable composition will generally contain a pharmaceutically acceptable amount of a subject oxazolidinone compound as a soluble salt dissolved in a liquid carrier such as water for injection to form a suitably buffered isotonic solution. Such a solution is stated

25 therein generally to contain a subject oxazolidinone compound dissolved in the carrier in an amount sufficient to provide an injectable concentration of about 1 to about 400 mg/ml of solution.

Many oxazolidinone compounds useful as antibiotics do not form, or do not readily form, salts. For these compounds, and where for any reason it is preferred not

to provide the antibiotic in salt form, it is generally difficult to formulate the antibiotic as a solution in a pharmaceutically acceptable liquid carrier, particularly an aqueous carrier. Most such compounds have relatively low solubility in water; in the case of linezolid, for example, the solubility at ambient temperature is less than 3 mg/ml and the practical limit of concentration in aqueous solution is about 2 mg/ml.

Particularly where parenteral or oral administration of an oxazolidinone antibiotic drug is contemplated, it is desired to achieve systemic concentrations of the drug in the bloodstream above a minimum inhibitory concentration for 90% of target organisms (MIC₉₀). It will readily be understood that it is difficult to achieve such concentrations by administration of a relatively small volume of a composition wherein the drug is present in dissolved form, unless the composition has a relatively high drug loading, and in particular a drug loading substantially above the limit of solubility in water of most oxazolidinone antibiotics not in the form of a salt.

A need therefore exists for a solution composition of an oxazolidinone antibiotic drug having a drug loading substantially in excess of the practical limit of solubility of the drug in water. A particular need exists for a parenterally deliverable solution composition of an oxazolidinone antibiotic drug having a relatively high concentration of the drug.

These and other needs will be seen to be met by the invention now described.

SUMMARY OF THE INVENTION

The present invention provides a pharmaceutical composition suitable for therapeutic or prophylactic administration to a subject having or at risk of infective disease, the composition comprising an aqueous carrier having in solution therein (a) an oxazolidinone antimicrobial drug in a therapeutically or prophylactically effective drug concentration that is above the practical limit of solubility of the drug in a substantially isotonic aqueous solution at a physiologically compatible pH, *e.g.*, pH 4.8, and (b) a pharmaceutically acceptable cyclodextrin compound in a concentration sufficient to maintain the drug in solution at such a drug concentration. Preferably the drug concentration is in a range of about 3 to about 100 mg/ml.

The term "suitable for therapeutic or prophylactic administration" in the present context encompasses compositions that are suitable for direct administration as formulated, and also compositions that are suitable for administration upon dilution

in an appropriate pharmaceutically acceptable liquid. Where the composition is intended for direct administration as formulated, the drug concentration is more preferably about 4 to about 40 mg/ml and most preferably about 5 to about 10 mg/ml. Where the composition is intended for dilution prior to administration, the drug concentration is more preferably about 10 to about 100 mg/ml and most preferably about 30 to about 100 mg/ml.

It is believed, without being bound by theory, that the enhanced solubility of the drug in a composition of the invention is due to association of at least a portion of the drug with the cyclodextrin. It is further believed that at least one mechanism by which the drug associates with the cyclodextrin compound to enhance solubility of the drug in an aqueous medium is through formation of an inclusion complex. Such complexes or conjugates are known in the art to form with a variety of drugs, and a number of advantages have been postulated for use of cyclodextrin-drug complexes in pharmacy. See for example review articles by Bekers *et al.* (1991) in Drug Development and Industrial Pharmacy, 17, 1503-1549; Szejtli (1994) in Medical Research Reviews, 14, 353-386; and Zhang & Rees (1999) in Expert Opinion on Therapeutic Patents, 9, 1697-1717.

Formulations of various drugs with various cyclodextrins have been proposed in the patent literature, including the patents and publications referenced below.

U.S. Patent No. 5,670,530 to Chen & Shishido discloses compositions comprising a rhodacyanine anti-cancer agent and a cyclodextrin.

U.S. Patent No. 5,756,546 to Pirotte *et al.* discloses compositions comprising nimesulide and a cyclodextrin.

U.S. Patent No. 5,807,895 to Stratton *et al.* discloses compositions comprising a prostaglandin and a cyclodextrin.

U.S. Patent No. 5,824,668 to Rubinfeld *et al.* discloses compositions comprising a 5β steroid drug and a cyclodextrin.

International Patent Publication No. WO 96/32135 discloses compositions comprising propofol and a cyclodextrin.

International Patent Publication No. WO 96/38175 discloses compositions comprising an antiulcerative benzimidazole compound and a branched cyclodextrin-carboxylic acid.

International Patent Publication No. WO 97/39770 discloses compositions comprising a thrombin inhibitor and a cyclodextrin.

International Patent Publication No. WO 98/37884 discloses compositions comprising a 3,4-diarylchroman compound and a cyclodextrin.

5 International Patent Publication No. WO 98/55148 discloses compositions comprising a sparingly water-soluble drug, a cyclodextrin, a water-soluble acid and a water-soluble organic polymer.

International Patent Publication No. WO 98/58677 discloses compositions comprising voriconazole and a cyclodextrin.

10 International Patent Publication No. WO 99/24073 discloses compositions comprising a taxoid such as paclitaxel or docetaxel and a cyclodextrin.

International Patent Publication No. WO 99/27932 discloses compositions comprising an antifungal compound of defined formula and a cyclodextrin.

However, the degree of enhancement of solubility achievable through
15 complexation with cyclodextrins of a particular drug or class of drugs is not generally predictable. Cyclodextrins are expensive excipients and in many cases the degree of enhancement of solubility, or other benefit obtained, has not economically justified the increased cost of a formulation arising from addition of a cyclodextrin. The present invention is based in part on the discovery that addition of a relatively modest
20 amount of a cyclodextrin compound increases the solubility of an oxazolidinone antibiotic drug to a surprising degree. This enhancement in solubility, among other benefits, makes it possible for the first time to deliver intravenously a therapeutically or prophylactically effective dose of the oxazolidinone in a volume small enough to be clinically acceptable and convenient, even for subjects intolerant of large volume
25 intravenous infusion because of hypertension, cardiac, renal and/or other problems. For example, a 600 mg dose of linezolid can, through use of a composition of the present invention, be delivered intravenously in a volume of 100 ml or less.

The term "pharmaceutically acceptable" in relation to a cyclodextrin or other excipient herein means having no persistent detrimental effect on the health of the
30 subject being treated. The pharmaceutical acceptability of a cyclodextrin depends, among other factors, on the particular cyclodextrin compound in question, on its concentration in the administered composition, and on the route of administration.

For example, use of β -cyclodextrin as an excipient in intravenous compositions is limited by hemolytic and nephrotoxic effects, but is generally non-toxic when administered orally.

5 The term "practical limit of solubility" in relation to a drug means the highest concentration at which the drug can be formulated in solution without risk of precipitation or crystallization of the drug during the normal range of manufacturing, packaging, storage, handling and use conditions. Typically the practical limit of solubility is considerably lower than the true solubility limit in a given aqueous medium, for example about 70% of the true solubility limit. Thus, illustratively, for a
10 drug having a true solubility limit in a given aqueous medium of 2.9 mg/ml, the practical limit of solubility is likely to be about 2 mg/ml.

Except where the context demands otherwise, use of the singular herein will be understood to embrace the plural. For example, by indicating above that a composition of the invention comprises "an oxazolidinone antibiotic drug" and "a
15 pharmaceutically acceptable cyclodextrin compound", it will be understood that the composition can contain one or more such drugs and one or more such cyclodextrin compounds.

The invention also provides a method of preparing a medicament for treating or preventing infective disease, using a composition as described above.

20 Also embraced by the present invention is a method of treating or preventing infective disease in a subject, the method comprising administration to the subject of a composition as described above in a therapeutically or prophylactically effective dose. Such administration can be oral, parenteral or topical, but is preferably parenteral and more preferably by intravenous injection or infusion.

25 The method of the invention is particularly useful where the infective disease arises through infection by one or more gram-positive bacteria, for example those of the genera *Staphylococcus* (e.g., *Staphylococcus aureus*, *Staphylococcus epidermidis*), *Streptococcus* (e.g., *Streptococcus viridans*, *Streptococcus pneumoniae*), *Enterococcus* (e.g., *Enterococcus faecalis*, *Enterococcus faecium*), *Bacillus*,
30 *Corynebacterium*, *Chlamydia* and *Neisseria*; anaerobic organisms, for example those of the genera *Bacteroides* and *Clostridia*; and acid-fast organisms, for example those of the genus *Mycobacterium*. The method of the invention is especially useful where


infection is by a strain of gram-positive bacteria that is resistant to one or more non-oxazolidinone antibiotics.

BRIEF DESCRIPTION OF THE DRAWING

Fig. 1 is a graphical representation of data from the study described in Example 6 herein, and demonstrates enhanced saturation solubility of oxazolidinone compounds in aqueous solutions containing hydroxypropyl- β -cyclodextrin (HP- β -CD).

DETAILED DESCRIPTION OF THE INVENTION

Any oxazolidinone antimicrobial drug, *i.e.*, one having an oxazolidinone moiety as part of its chemical structure, can be formulated with a cyclodextrin compound in accordance with the invention. Preferred oxazolidinones are compounds having formula (II):



where Ar is an optionally substituted aryl or heteroaryl group and R¹ is a group selected such that the compound of formula (II) falls within the scope of compounds generically or specifically disclosed in any of the following patents, each of which is individually incorporated herein by reference.

U.S. Patent No. 5,164,510 to Brickner.

U.S. Patent No. 5,231,188 to Brickner.

U.S. Patent No. 5,565,571 to Barbachyn & Brickner.

U.S. Patent No. 5,627,181 to Riedl *et al.*

U.S. Patent No. 5,652,238 to Barbachyn *et al.*


U.S. Patent No. 5,688,792 to Barbachyn *et al.*

U.S. Patent No. 5,698,574 to Riedl *et al.*

U.S. Patent No. 6,069,145 to Betts.

More preferably Ar is an optionally substituted 5- or 6-membered aryl or heteroaryl ring having 0 to 3 heteroatoms selected from nitrogen, oxygen and sulfur.

Still more preferably Ar is a group



where X is O, S, SO, SO₂, SNR⁴, S(O)NR⁴, NR⁴ or NC(O)CH₂OR⁴, where R⁴ is selected from hydrogen, R⁵ and -C(O)R⁵ groups where R⁵ is C₁₋₈ hydrocarbyl optionally substituted with one or more hydroxy, fluorine or chlorine groups; R⁶ and R⁷ are independently selected from hydrogen, methyl and cyano groups; and R⁸ and R⁹ are independently selected from hydrogen, fluorine and chlorine atoms. Most preferably R⁶ and R⁷ are hydrogen, and one of R⁸ and R⁹ is fluorine and the other of R⁸ and R⁹ is hydrogen.

Preferably R¹ is a group -(CH₂)_nN(R²)COR³ where n is 1 to 3, and R² and R³ are independently selected from hydrogen and C₁₋₈ hydrocarbyl optionally substituted with one or more hydroxy, fluorine or chlorine groups.

Examples of preferred oxazolidinones are compounds selected from linezolid, N-((5S)-3-(3-fluoro-4-(4-(2-fluoroethyl)-3-oxopiperazin-1-yl)phenyl)-2-oxooxazolidin-5-ylmethyl)acetamide, (S)-N-[[3-[5-(3-pyridyl)thiophen-2-yl]-2-oxo-5-oxazolidinyl]methyl]acetamide, (S)-N-[[3-[5-(4-pyridyl)pyrid-2-yl]-2-oxo-5-oxazolidinyl]methyl]acetamide hydrochloride and N-[[[(5S)-3-[4-(1,1-dioxido-4-thiomorpholinyl)-3,5-difluorophenyl]-2-oxo-5-oxazolidinyl]methyl]acetamide. In one especially preferred embodiment, the oxazolidinone is linezolid. In another especially preferred embodiment, the oxazolidinone is N-[[[(5S)-3-[4-(1,1-dioxido-4-thiomorpholinyl)-3,5-difluorophenyl]-2-oxo-5-oxazolidinyl]methyl]acetamide.

The invention is illustrated herein with particular reference to linezolid, and it will be understood that any other oxazolidinone antimicrobial drug can, if desired, be substituted in whole or in part for linezolid, with appropriate adjustment in concentration and dosage ranges, in the compositions and methods herein described.

Oxazolidinone compounds used in compositions of the invention can be prepared by a process known *per se*, in the case of linezolid, for example, by processes described in the following patents, each of which is individually incorporated herein by reference.

Above-cited U.S. Patent No. 5,688,792.

U.S. Patent No. 5,837,870 to Barbachyn *et al.*

International Patent Publication No. WO 99/24393.

Other oxazolidinone compounds can be prepared by processes known *per se*, including processes set forth in patent publications disclosing such drugs.

5 Linezolid is usefully present in a composition of the invention at a concentration of about 3 mg/ml to as high a concentration as is practically enabled by the cyclodextrin present therewith, for example about 100 mg/ml. Preferably in a composition intended for direct administration as formulated, the concentration of linezolid is about 4 to about 40 mg/ml, more preferably about 5 to about 10 mg/ml, for
10 example about 6 mg/ml. Preferably in a composition intended for dilution in a pharmaceutically acceptable liquid prior to administration, the concentration of linezolid is about 10 to about 100 mg/ml, more preferably about 30 to about 100 mg/ml, for example about 60 mg/ml. Useful concentrations of other oxazolidinone drugs are those that are therapeutically equivalent to the linezolid concentration ranges
15 given immediately above.

 The cyclodextrin compound with which the oxazolidinone antibiotic drug is formulated according to the present invention is preferably selected from α -cyclodextrin, β -cyclodextrin, γ -cyclodextrin, alkylcyclodextrins (*e.g.*, methyl- β -cyclodextrin, dimethyl- β -cyclodextrin, diethyl- β -cyclodextrin),
20 hydroxyalkylcyclodextrins (*e.g.*, hydroxyethyl- β -cyclodextrin, hydroxypropyl- β -cyclodextrin), carboxyalkylcyclodextrins (*e.g.*, carboxymethyl- β -cyclodextrin) and sulfoalkylether cyclodextrins (*e.g.*, sulfobutylether- β -cyclodextrin). More preferred are hydroxyalkyl- β -cyclodextrins and sulfoalkylether- β -cyclodextrins; still more preferred are hydroxypropyl- β -cyclodextrin and sulfobutylether- β -cyclodextrin, with
25 sulfobutylether- β -cyclodextrin being presently the most preferred.

 If desired, complexation of an oxazolidinone antibiotic drug by a cyclodextrin can be increased by addition of a water-soluble polymer such as carboxymethylcellulose or a salt thereof, hydroxypropylmethylcellulose or polyvinylpyrrolidone, as described by Loftsson (1998), Pharmazie, 53, 733-740.

30 The cyclodextrin is present at a concentration effective to enhance the solubility of the oxazolidinone, for example at a concentration of about 1 to about 500 mg/ml. In practice and in view of the high cost of cyclodextrins, the amount of the

cyclodextrin present in a composition of the invention is preferably only slightly greater, for example no more than about 50% greater, than a minimum amount required to maintain the oxazolidinone in solution at the desired oxazolidinone concentration.

5 Typically, where the composition is intended for direct administration as formulated, suitable concentrations of cyclodextrin will be found in a range from about 5 to about 200 mg/ml, more commonly about 10 to about 100 mg/ml, and most commonly about 20 to about 80 mg/ml. Where the composition is intended for dilution prior to administration, the concentration of cyclodextrin can be significantly
10 higher, for example about 100 to about 500 mg/ml.

 One or more pharmaceutically acceptable pH adjusting agents and/or buffering agents can be included in a composition of the invention, including acids such as acetic, boric, citric, lactic, phosphoric and hydrochloric acids; bases such as sodium hydroxide, sodium phosphate, sodium borate, sodium citrate, sodium acetate, sodium
15 lactate and tris-hydroxymethylaminomethane; and buffers such as citrate/dextrose, citrate/phosphate, sodium bicarbonate and ammonium chloride. Such acids, bases and buffers are included in an amount required to maintain pH of the composition in a physiologically acceptable range, particularly where the composition is intended for parenteral delivery.

20 One or more pharmaceutically acceptable salts or other solutes can be included in the composition in an amount required to bring osmolality of the composition into a physiologically acceptable range, particularly where the composition is intended for parenteral delivery. Such salts include those having sodium, potassium or ammonium cations and chloride, citrate, ascorbate, borate, phosphate, bicarbonate, sulfate,
25 thiosulfate or bisulfite anions; preferred salts include sodium chloride, potassium chloride, sodium thiosulfate, sodium bisulfite and ammonium sulfate, with sodium chloride being especially preferred. Other solutes suitable for adjustment of osmolality include sugars, for example dextrose.

 Accordingly, a particular embodiment of the invention is a composition as
30 described hereinabove, further comprising a buffering agent and/or an agent for adjusting osmolality in amounts whereby the solution is substantially isotonic and has a physiologically acceptable pH.

Other pharmaceutically acceptable excipients can also be included as desired in compositions of the invention, having functions conventional in the art and in amounts consistent with those functions. For example, a water-soluble organic solvent can be included if desired, as disclosed in U.S. Patent No. 5,486,508 to Nishida *et al.*, which contemplates a composition suitable for injection comprising a slightly water-soluble drug, a cyclodextrin and a water-soluble organic solvent.

Compositions of the present invention can be prepared by processes known in the art, including by simple admixture, with agitation as appropriate, of the ingredients. Preferably an aqueous solution of the cyclodextrin compound is first prepared, and the oxazolidinone in finely divided solid particulate form is added to that solution with agitation until it is fully dissolved. Where it is desired to prepare a buffered isotonic solution, for example for intravenous infusion, buffering agents and agents for adjustment of osmolality can be added at any stage but are preferably present in solution with the cyclodextrin compound before addition of the oxazolidinone. Processes for preparing a composition of the invention, particularly one intended for parenteral use, are preferably conducted so as to provide a sterile product.

Compositions of the invention intended for parenteral administration are generally suitable for packaging and dispensing in conventional intravenous delivery bags and apparatus.

A contemplated composition can be dried, for example by spray drying, to form a reconstitutable powder. The powder can be dissolved in sterile water to reconstitute a parenterally deliverable composition as herein described.

In a method of the invention for treating or preventing infective disease, a composition as described above in a therapeutically or prophylactically effective daily dose is administered to a subject in need thereof. Such administration can be oral, parenteral or topical, but is preferably parenteral and more preferably by intravenous injection or infusion.

In a particular embodiment of the invention, a method for treating or preventing infective disease comprises (a) diluting a composition as described herein in a pharmaceutically acceptable liquid to form a diluted composition suitable for direct administration, and (b) administering the diluted composition in a

therapeutically or prophylactically effective daily dose to a subject in need thereof. Preferably such administration is parenteral and the liquid in which the composition is diluted is a parenterally acceptable aqueous carrier, for example saline or a substantially isotonic buffered aqueous solution having a physiologically compatible pH.

As indicated above, a method of the invention is particularly useful where the infective disease arises through infection by one or more gram-positive bacteria. Where broader-spectrum antibacterial activity, extending to gram-negative bacteria, is required, a second antimicrobial drug can be administered in co-therapy, including for example coformulation, with the present composition. The second antimicrobial drug is selected to be effective against target gram-negative bacteria. Such co-therapy and coformulation are embodiments of the present invention.

The second antimicrobial drug can illustratively be selected from aminoglycosides, cephalosporins, diaminopyridines, fluoroquinolones, sulfonamides and tetracyclines. Among particular antimicrobial drugs of these and other classes, each of the following may illustratively be useful as the second antimicrobial drug according to an embodiment of the present invention: amikacin, cefixime, cefoperazone, cefotaxime, ceftazidime, ceftizoxime, ceftriaxone, chloramphenicol, ciprofloxacin, clindamycin, colistin, domeclocycline, doxycycline, gentamicin, mafenide, methacycline, minocycline, neomycin, norfloxacin, ofloxacin, oxytetracycline, polymyxin B, pyrimethamine, silver sulfadiazine, sulfacetamide, sulfisoxazole, tetracycline, tobramycin and trimethoprim.

The present invention also encompasses therapeutic and prophylactic methods involving administration of an antibacterial composition as described herein in co-therapy, including for example coformulation, with one or more drugs other than antibacterial drugs.

Therapeutic and prophylactic methods of the invention are useful for any subject in need thereof. The subject is preferably warm-blooded, more preferably mammalian, and most preferably human. However, a particular embodiment of the invention is a veterinary method of treating a non-human subject, for example a domestic, farm or zoo animal, having or at risk of infective disease, with a composition of the invention.

An appropriate dosage, frequency and duration of administration, *i.e.*, treatment regimen, to be used in any particular situation will be readily determined by one of skill in the art without undue experimentation, and will depend, among other factors, on the particular oxazolidinone compound(s) present in the composition, on the particular infective disease or condition to be treated or prevented, on the age, weight and general physical condition of the subject, and on other medication being administered to the subject. It is preferred that response to treatment according to the present method be monitored and the treatment regimen be adjusted if necessary in light of such monitoring.

Where the oxazolidinone is linezolid, a daily dose for a human subject will typically be about 100 mg to about 1000 mg, more typically about 250 mg to about 750 mg, for example about 600 mg of linezolid, administered in a composition of the invention. For other oxazolidinones, a daily dose that is therapeutically equivalent to the above dose ranges for linezolid can be administered.

EXAMPLES

The following Examples illustrate aspects of the present invention but are not to be construed as limitations.

Example 1

A study was conducted to examine solubility of linezolid in an aqueous system containing sulfobutylether- β -cyclodextrin (SB- β -CD).

Aqueous solutions of SB- β -CD at concentrations of 10, 50, 100, 150, 250 and 500 mg/ml were prepared. Excess linezolid was added to each solution. The solutions were stirred for 24 h at 25°C and were then filtered using 0.2 μ m Gelman Acrodisc filter units and assayed for linezolid by HPLC.

Saturation solubility of linezolid in pure water at pH 7 was determined separately to be 2.9 ± 0.1 mg/ml. Saturation solubility of linezolid in aqueous SB- β -CD solutions was determined as shown in Table 1.

Table 1. Saturation solubility of linezolid in SB- β -CD solutions

SB-β-CD concentration (mg/ml)	Solubility of linezolid (mg/ml)
10	4.3
50	9.5
100	15.9
150	22.1
250	33.4
500	59.9

Example 2

A buffered isotonic solution was prepared at pH 4.5, 283 mOsm/kg using a 10mM citrate buffer solution (0.851 mg/ml citric acid, 1.638 mg/ml sodium citrate) containing 23 mg/ml dextrose and 50 mg/ml SB- β -CD. To 20 ml of this solution was added 100 mg linezolid with heating and stirring until the linezolid was completely dissolved. The resulting solution contained approximately 5 mg/ml linezolid.

Example 3

A buffered isotonic solution was prepared at pH 4.5, 285 mOsm/kg using a 10mM citrate buffer solution (0.851 mg/ml citric acid, 1.638 mg/ml sodium citrate) containing 100 mg/ml SB- β -CD. To 20 ml of this solution was added 200 mg linezolid with heating and stirring until the linezolid was completely dissolved. The resulting solution contained approximately 10 mg/ml linezolid.

Example 4

A buffered isotonic solution was prepared at pH 4.5, 289 mOsm/kg using a 10mM citrate buffer solution (0.851 mg/ml citric acid, 1.638 mg/ml sodium citrate) containing 100 mg/ml SB- β -CD. To 100 ml of this solution was added 800 mg linezolid with heating to about 60°C and stirring until the linezolid was completely dissolved. The resulting solution contained approximately 8 mg/ml linezolid.

The solution was filtered using 0.2 μ m Nalgene filterware, and 10 ml of filtered solution was filled into each of ten Kimble-Warsaw Type I glass bottles stoppered with Daikyo 777 stoppers. Eight of the bottles were placed in a 25°C constant temperature cabinet and two of the bottles were placed in a laboratory refrigerator at 4-6°C.

No precipitation or color changes were evident after storage for two months under these conditions.

Example 5

Four formulations X, A, B and C suitable for direct intravenous (IV) infusion, and one formulation D suitable for dilution with a standard IV diluent (*e.g.*, saline or 5% dextrose) are prepared, each containing a 600 mg dose of linezolid. Formulation X is included for comparative purposes, containing no cyclodextrin. Compositions of formulation X and formulations A-D are shown in Table 2.

Table 2. Compositions of formulations X and A-D (mg/ml)

	X	A	B	C	D
Linezolid	2	8	10	8	60
Sulfobutylether- β -cyclodextrin	-	100	100	50	500
Sodium citrate dihydrate	1.638	1.638	1.638	1.638	1.638
Citric acid monohydrate granular	0.851	0.851	0.851	0.851	0.851
Dextrose anhydrous granular	*	-	-	23	-
Water for injection USP	q.s.	q.s.	q.s.	q.s.	q.s.
Total volume (ml) containing 600 mg dose	300	75	60	75	10

* concentration required to provide isotonic solution in absence of cyclodextrin

q.s. = quantity sufficient to make up to total volume shown

The very substantial reduction in volume of IV solution providing a 600 mg dose of linezolid, permitted by addition of SB- β -CD, is clearly seen in Table 2 above.

Example 6

A study was conducted to examine solubility of three oxazolidinone compounds, herein denoted Compound 1, Compound 2 and Compound 3, in an aqueous system containing hydroxypropyl- β -cyclodextrin (HP- β -CD).

Compound 1 is (S)-N-[[3-[3-fluoro-4-(4-(hydroxyacetyl)-1-piperazinyl)phenyl]-2-oxo-5-oxazolidinyl]methyl]acetamide.

Compound 2 is (S)-N-[[3-[3-fluoro-4-(4-morpholinyl)phenyl]-2-oxo-5-oxazolidinyl]methyl]acetamide (linezolid).

Compound 3 is (S)-N-[[3-[3-fluoro-4-(1,1-dioxothiomorpholin-4-yl)phenyl]-2-oxo-5-oxazolidinyl]methyl]acetamide.

Aqueous solutions of HP- β -CD at concentrations of 0, 60, 100, 200, 300 and 400 mg/ml were prepared. Compound 1, 2 or 3 in excess amount was added to each solution. The solutions were stirred for 48 h at 37°C and were then filtered and assayed by HPLC to provide a measure of saturation solubility of Compounds 1, 2 and

3 in each HP- β -CD solution.

The saturation solubilities are shown in graphical form in Fig. 1. Saturation solubility of each oxazolidinone compound was found to be linearly related to HP- β -CD concentration.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000
1001
1002
1003
1004
1005
1006
1007
1008
1009
1010
1011
1012
1013
1014
1015
1016
1017
1018
1019
1020
1021
1022
1023
1024
1025
1026
1027
1028
1029
1030
1031
1032
1033
1034
1035
1036
1037
1038
1039
1040
1041
1042
1043
1044
1045
1046
1047
1048
1049
1050
1051
1052
1053
1054
1055
1056
1057
1058
1059
1060
1061
1062
1063
1064
1065
1066
1067
1068
1069
1070
1071
1072
1073
1074
1075
1076
1077
1078
1079
1080
1081
1082
1083
1084
1085
1086
1087
1088
1089
1090
1091
1092
1093
1094
1095
1096
1097
1098
1099
1100
1101
1102
1103
1104
1105
1106
1107
1108
1109
1110
1111
1112
1113
1114
1115
1116
1117
1118
1119
1120
1121
1122
1123
1124
1125
1126
1127
1128
1129
1130
1131
1132
1133
1134
1135
1136
1137
1138
1139
1140
1141
1142
1143
1144
1145
1146
1147
1148
1149
1150
1151
1152
1153
1154
1155
1156
1157
1158
1159
1160
1161
1162
1163
1164
1165
1166
1167
1168
1169
1170
1171
1172
1173
1174
1175
1176
1177
1178
1179
1180
1181
1182
1183
1184
1185
1186
1187
1188
1189
1190
1191
1192
1193
1194
1195
1196
1197
1198
1199
1200
1201
1202
1203
1204
1205
1206
1207
1208
1209
1210
1211
1212
1213
1214
1215
1216
1217
1218
1219
1220
1221
1222
1223
1224
1225
1226
1227
1228
1229
1230
1231
1232
1233
1234
1235
1236
1237
1238
1239
1240
1241
1242
1243
1244
1245
1246
1247
1248
1249
1250
1251
1252
1253
1254
1255
1256
1257
1258
1259
1260
1261
1262
1263
1264
1265
1266
1267
1268
1269
1270
1271
1272
1273
1274
1275
1276
1277
1278
1279
1280
1281
1282
1283
1284
1285
1286
1287
1288
1289
1290
1291
1292
1293
1294
1295
1296
1297
1298
1299
1300
1301
1302
1303
1304
1305
1306
1307
1308
1309
1310
1311
1312
1313
1314
1315
1316
1317
1318
1319
1320
1321
1322
1323
1324
1325
1326
1327
1328
1329
1330
1331
1332
1333
1334
1335
1336
1337
1338
1339
1340
1341
1342
1343
1344
1345
1346
1347
1348
1349
1350
1351
1352
1353
1354
1355
1356
1357
1358
1359
1360
1361
1362
1363
1364
1365
1366
1367
1368
1369
1370
1371
1372
1373
1374
1375
1376
1377
1378
1379
1380
1381
1382
1383
1384
1385
1386
1387
1388
1389
1390
1391
1392
1393
1394
1395
1396
1397
1398
1399
1400
1401
1402
1403
1404
1405
1406
1407
1408
1409
1410
1411
1412
1413
1414
1415
1416
1417
1418
1419
1420
1421
1422
1423
1424
1425
1426
1427
1428
1429
1430
1431
1432
1433
1434
1435
1436
1437
1438
1439
1440
1441
1442
1443
1444
1445
1446
1447
1448
1449
1450
1451
1452
1453
1454
1455
1456
1457
1458
1459
1460
1461
1462
1463
1464
1465
1466
1467
1468
1469
1470
1471
1472
1473
1474
1475
1476
1477
1478
1479
1480
1481
1482
1483
1484
1485
1486
1487
1488
1489
1490
1491
1492
1493
1494
1495
1496
1497
1498
1499
1500
1501
1502
1503
1504
1505
1506
1507
1508
1509
1510
1511
1512
1513
1514
1515
1516
1517
1518
1519
1520
1521
1522
1523
1524
1525
1526
1527
1528
1529
1530
1531
1532
1533
1534
1535
1536
1537
1538
1539
1540
1541
1542
1543
1544
1545
1546
1547
1548
1549
1550
1551
1552
1553
1554
1555
1556
1557
1558
1559
1560
1561
1562
1563
1564
1565
1566
1567
1568
1569
1570
1571
1572
1573
1574
1575
1576
1577
1578
1579
1580
1581
1582
1583
1584
1585
1586
1587
1588
1589
1590
1591
1592
1593
1594
1595
1596
1597
1598
1599
1600
1601
1602
1603
1604
1605
1606
1607
1608
1609
1610
1611
1612
1613
1614
1615
1616
1617
1618
1619
1620
1621
1622
1623
1624
1625
1626
1627
1628
1629
1630
1631
1632
1633
1634
1635
1636
1637
1638
1639
1640
1641
1642
1643
1644
1645
1646
1647
1648
1649
1650
1651
1652
1653
1654
1655
1656
1657
1658
1659
1660
1661
1662
1663
1664
1665
1666
1667
1668
1669
1670
1671
1672
1673
1674
1675
1676
1677
1678
1679
1680
1681
1682
1683
1684
1685
1686
1687
1688
1689
1690
1691
1692
1693
1694
1695
1696
1697
1698
1699
1700
1701
1702
1703
1704
1705
1706
1707
1708
1709
1710
1711
1712
1713
1714
1715
1716
1717
1718
1719
1720
1721
1722
1723
1724
1725
1726
1727
1728
1729
1730
1731
1732
1733
1734
1735
1736
1737
1738
1739
1740
1741
1742
1743
1744
1745
1746
1747
1748
1749
1750
1751
1752
1753
1754
1755
1756
1757
1758
1759
1760
1761
1762
1763
1764
1765
1766
1767
1768
1769
1770
1771
1772
1773
1774
1775
1776
1777
1778
1779
1780
1781
1782
1783
1784
1785
1786
1787
1788
1789
1790
1791
1792
1793
1794
1795
1796
1797
1798
1799
1800
1801
1802
1803
1804
1805
1806
1807
1808
1809
1810
1811
1812
1813
1814
1815
1816
1817
1818
1819
1820
1821
1822
1823
1824
1825
1826
1827
1828
1829
1830
1831
1832
1833
1834
1835
1836
1837
1838
1839
1840
1841
1842
1843
1844
1845
1846
1847
1848
1849
1850
1851
1852
1853
1854
1855
1856
1857
1858
1859
1860
1861
1862
1863
1864
1865
1866
1867
1868
1869
1870
1871
1872
1873
1874
1875
1876
1877
1878
1879
1880
1881
1882
1883
1884
1885
1886
1887
1888
1889
1890
1891
1892
1893
1894
1895
1896
1897
1898
1899
1900
1901
1902
1903
1904
1905
1906
1907
1908
1909
1910
1911
1912
1913
1914
1915
1916
1917
1918
1919
1920
1921
1922
1923
1924
1925
1926
1927
1928
1929
1930
1931
1932
1933
1934
1935
1936
1937
1938
1939
1940
1941
1942
1943
1944
1945
1946
1947
1948
1949
1950
1951
1952
1953
1954
1955
1956
1957
1958
1959
1960
1961
1962
1963
1964
1965
1966
1967
1968
1969
1970
1971
1972
1973
1974
1975
1976
1977
1978
1979
1980
1981
1982
1983
1984
1985
1986
1987
1988
1989
1990
1991
1992
1993
1994
1995
1996
1997
1998
1999
2000
2001
2002
2003
2004
2005
2006
2007
2008
2009
2010
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022
2023
2024
2025
2026
2027
2028
2029
2030
2031
2032
2033
2034
2035
2036
2037
2038
2039
2040
2041
2042
2043
2044
2045
2046
2047
2048
2049
2050
2051
2052
2053
2054
2055
2056
2057
2058
2059
2060
2061
2062
2063
2064
2065
2066
2067
2068
2069
2070
2071
2072
2073
2074
2075
2076
2077
2078
2079
2080
2081
2082
2083
2084
2085
2086
2087
2088
2089
2090
2091
2092
2093
2094
2095
2096
2097
2098
2099
2100
2101
2102
2103
2104
2105
2106
2107
2108
2109
2110
2111
2112
2113
2114
2115
2116
2117
2118
2119
2120
2121
2122
2123
2124
2125
2126
2127
2128
2129
2130
2131
2132
2133
2134
2135
2136
2137
2138
2139
2140
2141
2142
2143
2144
2145
2146
2147
2148
2149
2150
2151
2152
2153
2154
2155
2156
2157
2158
2159
2160
2161
2162
2163
2164
2165
2166
2167
2168
2169
2170
2171
2172
2173
2174
2175
2176
2177
2178
2179
2180
2181
2182
2183
2184
2185
2186
2187
2188
2189
2190
2191
2192
2193
2194
2195
2196
2197
2198
2199
2200
2201
2202
2203
2204
2205
2206
2207
2208
2209
2210
2211
2212
2213
2214
2215
2216
221